

Linear Regression: Analytic Solution*(find and work with a partner)*

In the case of *multiple* linear regression, each example has p features. We still have an associated response (output) variable y for each example. We typically add a “fake 1” to the features of each example, so that $\mathbf{x} = [1 \ x_1 \ x_2 \ \cdots \ x_p]^T$. Our model is now the dot product between the weight vector \mathbf{w} and the features:

$$h_{\mathbf{w}}(\mathbf{x}) = w_0 + w_1x_1 + \cdots + w_px_p = \mathbf{w} \cdot \mathbf{x}$$

If we consider \mathbf{X} to be the entire matrix of features (with dimensions $n \times (p + 1)$) then we can reframe the linear regression problem in terms of vectors and matrices. In this case the analytic solution for the weight vector is:

$$\hat{\mathbf{w}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

1. *Warmup.* Given matrices \mathbf{A} and \mathbf{B} below, compute both \mathbf{AB} and \mathbf{BA} . Is matrix multiplication commutative?

$$\mathbf{A} = \begin{bmatrix} 1 & -1 \\ 3 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 0 & 2 \\ -4 & 1 \end{bmatrix}$$

2. Going back to our small example from the last handout, we will use this matrix/vector form to verify our solution for \mathbf{w} (even though $p = 1$ we can still use the multiple linear regression derivation). We again have the data: $(x_1, y_1) = (1, 0)$ and $(x_2, y_2) = (0, 1)$. First, rewrite \mathbf{X} with the added column of 1's. Then write \vec{y} as a vector.
3. Now use the matrix/vector analytic solution to verify the weight vector \mathbf{w} .